

Do you have a healthy heart?

Almost **1** out of
3 people die
from cardiovascular disease.

In fact, cardiovascular disease is responsible for 31% of deaths among men and 33% among women. Cardiovascular disease is health enemy number 1 and is the principal cause of disability in Canada.

Cardiovascular disease affects the heart (cardio) and blood vessels (vascular), triggering angina or heart attacks (myocardial infarct), as well as strokes (cerebral vascular accidents) that can result in major paralysis.

The health of your heart is in your hands.

Take the
1 test

2 Assess your
risk level

3 Take
action



Do you have a healthy heart?

- This test is designed for adults. Pregnant women and breastfeeding mothers should wait until the baby has been weaned to take the test.
- For each question, circle the score that best corresponds to your situation and transfer it to the rectangle on the right.
- For questions 7, 8 and 9, a space has been provided to write your results. These results can be obtained from your doctor or as the result of a screening test.



A

1 Age

I am:

Under 40	0
40 to 49	2
50 to 59	5
60 to 69	8
70 and over	10

2 Gender

I am a man	8
I am a non-menopausal woman *	0
I am a menopausal woman * (for 10 years or less)	4
I am a menopausal woman * (for over 10 years)	8

*A woman is menopausal if menstruation has stopped for at least one year. She will remain menopausal for the remainder of her life.

3 Heredity

Is there a history of cardiovascular disease (angina, heart attack, coronary bypass surgery, paralysis) in your family? (birth father, mother, brothers or sisters)

No	0
Yes: who and at what age?	
• Father or brother at age 55 or under	8
• Father or brother over age 55	4
• Mother or sister at age 65 or under	8
• Mother or sister over age 65	4

If two answers are possible, choose the one with the highest score.

YOUR A TOTAL

Add up your scores for questions 1 to 3

B

4 Physical activity

In general, how many days per week are you physically active for at least 30 minutes? * (walking, dancing, sports, workout, etc.)

5 days or more per week	0
3 or 4 days per week:	
• high intensity (ex.: jogging, cycling at over 20 to 25 km/h)	0
• light or moderate intensity	3
1 or 2 days per week	6
Less than once per week	8

* Does not have to be a continuous 30 minutes.

5 Tobacco

I don't smoke	0
I don't smoke but I am regularly exposed to second-hand smoke	3
I only smoke cigars or a pipe	3
I smoke 10 cigarettes or less per day	4
I smoke 11 to 20 cigarettes per day	7
I smoke more than 20 cigarettes per day	10

For women who smoke and take birth control pills:

If you are under age 35, add	4
If you are aged 35 or over, add	10

Important: This test has been designed in light of the latest scientific knowledge. It is a tool to raise awareness of your current level of risk for cardiovascular disease. It also highlights the important role you can play to reduce that risk. It does not constitute a diagnosis and is no substitute for a consultation with your physician. Although certain additional risk factors of cardiovascular disease are receiving greater attention, they were not selected for inclusion in this test because their impact on cardiovascular disease risk levels has not yet been clearly established.



6 Weight and waist circumference

My height: cm or ft. in.
 My weight: kg or lbs

Refer to the weight chart on page 11 to identify your current weight zone:

Zone 1 - underweight	0
Zone 2 - normal weight	0
Zone 3 - overweight	4
Zone 4 - obesity	6

Measure your waist circumference at your navel.

Women:	
• Less than 88 cm (35 in)	0
• 88 cm (35 in) or more	4
Men:	
• Less than 102 cm (40 in)	0
• 102 cm (40 in) or more	4

7 Diabetes

My blood sugar level: mmol/L

Have you ever had a blood test to determine if you have diabetes?

No, and there are no diabetics in my family (birth father, mother, brothers or sisters)	4
No, but there are diabetics in my family (birth father, mother, brothers or sisters)	8
Yes, and I am not diabetic	0
Yes, and I was told I am at risk of becoming diabetic (slightly elevated blood sugar level: pre-diabetic or glucose intolerant)	15
Yes, and I am diabetic	30

8 Blood pressure

My reading: / mm Hg

I don't know what my blood pressure is	4
My blood pressure is:	
• Normal: Less than 130/85	0
• At the limit: 130/85 to 139/89	2
• Slightly elevated: 140/90 to 159/99	6
• High: 160/100 or over	10

9 Cholesterol and triglycerides

A) Total cholesterol

My result: mmol/L

I don't know my total blood cholesterol level	2
My total cholesterol level is:	
• Normal: under 5.2 mmol/L	0
• Slightly elevated: 5.2 to 6.1 mmol/L	3
• Elevated: 6.2 mmol/L or higher	4

B) HDL cholesterol (good cholesterol)

My result: mmol/L

I don't know my HDL cholesterol level	1
My HDL cholesterol level is:	
• Normal: 1.0 mmol/L or more	0
• Too low: under 1.0 mmol/L	4

C) LDL cholesterol (bad cholesterol)

My result: mmol/L

I don't know my LDL cholesterol level	1
My LDL cholesterol level is:	
• Normal: under 3.4 mmol/L	0
• High: 3.4 mmol/L or higher	4

D) Triglycerides

My result: mmol/L

I don't know my triglyceride level	1
My triglyceride level is:	
• Normal: under 1.7 mmol/L	0
• High: 1.7 mmol/L or higher	4

YOUR B TOTAL

Add up your scores for questions 4 to 9

Copy

YOUR A TOTAL

YOUR B TOTAL

on page 4.

There are certain factors that put you at a greater risk of developing cardiovascular disease. The more risk factors you have, the higher your risk.

YOUR A TOTAL

is your score for the **3 non-modifiable risk factors**. It is recognized that risk increases with age, that women are affected later than men, i.e. after the onset of menopause, and that certain families are more prone to cardiovascular disease.



YOUR B TOTAL

is your score for the **6 modifiable risk factors**. Your lifestyle choices can play an important role.



YOUR A+B TOTAL

YOUR A + B TOTAL = YOUR RISK LEVEL

0 to 11

very low risk

12 to 21

low risk

If your level of risk is very low or low:

Congratulations! To maintain this level throughout your life, be sure to keep the modifiable risk factors under control.

22 to 30

average risk

If your level of risk is average:

You can improve your score through changes to your lifestyle. Look at how your points are distributed among the modifiable risk factors and take action to increase your chances of staying healthy.

31 to 42

high risk

43 or more

very high risk

If your level of risk is high or very high:

We suggest you talk to your physician. In the meantime, look at your highest scores for modifiable factors, and start thinking of ways to lower your risk level.

BEWARE!

- If you have already had a heart attack, stroke or episodes of angina, your risk level is considered very high, whatever your results to this questionnaire. The scores show which factors to control in order to lower the risk of a new attack.
- If you don't know your blood pressure, your cholesterol levels, your triglyceride or blood sugar levels, we recommend you see your physician, especially if your risk level is high or very high.

No **time?** Too much **trouble?**

It's not always easy to take action. Have you ever asked yourself what's stopping you from making a priority of your heart's health? There could be several reasons. Identifying them is a step in the right direction.

Maybe you think you'll have to give up certain pleasures when you make a change for the better. Try looking at it a different way. Enjoying a better quality of life, having more energy and being in better shape are pleasures too, readily accessible and for a long time!

Your **heart** is in your **hands**

Age, gender and heredity are risk factors that cannot be modified. **The good news is there are six other factors that are largely under our control.**

4 WINNING STRATEGIES

EAT HEALTHY



**MAINTAIN
OR REDUCE
YOUR WEIGHT**

**BE ACTIVE 30 min.
ON MOST DAYS**

DO NOT SMOKE

**TO ACT ON THE
6 MODIFIABLE
RISK FACTORS
FOR CARDIOVASCULAR
DISEASE.**

1. physical inactivity
2. tobacco
3. excess weight
4. diabetes
5. high blood pressure
6. elevated cholesterol and/or triglyceride levels

Getting to understand the 6 modifiable risk factors

Physical inactivity *(Question 4)*

Inactive people run the same risk for cardiovascular disease they would if they smoked a pack of cigarettes a day.

Sports are certainly a good way to be more active but learn to take advantage of other activities that encourage you to move, such as walking, cycling, taking the stairs or playing outside with the kids.

What's important is to be active at least thirty minutes most days of the week. If you have the chance to exercise longer and more vigorously, you will get even more benefit out of it.

CAUTION: If your test indicates a high or very high risk level, or if you already have heart problems, see your physician before undertaking an intensive exercise program.

ONE OF **4** WINNING STRATEGIES



Tobacco *(Question 5)*

Healthwise, not smoking is the best investment you can make and, as far as the heart is concerned, the risk of cardiovascular disease drops the moment you quit.

Among women, combining the contraceptive pill and cigarettes significantly increases the risk, particularly for those aged 35 and over. Living and working with smokers also increases the risk of health problems in non-smokers.

Although quitting is not easy, the good news is there are now many methods and resources available to help you quit. When you decide to quit smoking, good preparation and the use of devices such as nicotine patches, gum or lozenges, as well as varenicline or bupropion tablets can really boost your chances of success. **The only way to fail is to stop trying!**

ONE OF **4** WINNING STRATEGIES



Excess weight *(Question 6)*

Excess weight, especially around the waistline, increases the risks of cardiovascular disease. The weight chart on page 11 gives a range of acceptable weights according to height (zone 2). It's up to you to determine the weight that suits you best within that range. As for waist circumference, experts consider that it should be less than 102 cm (40 inches) for men and 88 cm (35 inches) for women.

Losing some weight is a good idea as long as you don't gain it back (and more!), which is generally the case with crash diets and miracle products. The best way to lose weight is to do it gradually, by watching what you eat and maintaining a good level of physical activity.

If you are significantly overweight, you don't need to reach your 'normal weight' category to benefit from positive health results. Losing 5 to 10% of your current weight often improves the control of diabetes, cholesterol and blood pressure. It is far better to set a realistic goal that you can re-assess once you've reached it. When your weight is normal, maintaining that weight over the years is the best goal to choose.

What about the scale? Even if your clothes might tell you that you have gained a little weight, the scale remains the best tool to measure your weight and to manage it appropriately. It is much easier to lose 1 or 2 kilograms over a few days than to have 10 or even 20 kilograms to lose at a given moment.

ONE OF **4** WINNING STRATEGIES





The three risk factors on this page are health problems in which heredity plays an important role. It is, however, possible to prevent, delay or control them by adopting the “4 winning strategies” on page 5, as well as by following the advice on healthy eating on pages 8 and 9. If you must take medication, take it as prescribed. Don't change it without consulting your physician or pharmacist.

Diabetes (Question 7)

The risk for cardiovascular disease is up to four times greater for a diabetic than a non diabetic. In the long run, diabetes can lead to other complications such as vision impairment and kidney problems.

Measuring the level of sugar in your blood (glycemia) makes it possible to detect diabetes. It is even more important to know what your level is if there are diabetics in your family or if you are overweight. Diabetes does not develop overnight: if your physician tells you that your blood sugar is slightly above normal, measures can be taken to delay the onset of diabetes.

High blood pressure (Question 8)

Hypertension works silently: very often, there are no symptoms. The situation must be corrected, however, to avoid complications to the heart, brain or kidney.

The only way to know your blood pressure is to have it checked. Since blood pressure can fluctuate, it's necessary to have readings taken at different times before hypertension can be diagnosed. Normal blood pressure is under 130/85 mmHg. In certain cases, your physician can consider a higher blood pressure reading as acceptable.

Elevated Cholesterol (Question 9) and/or triglyceride levels

Cholesterol is a fat found in the blood. Your body needs it to produce hormones, bile, and cellular membranes. The problem starts when there's too much of it, which is the case for almost every other Canadian.

To travel in the blood, cholesterol needs carriers. There are two main kinds: the good carriers (HDL) and the bad carriers (LDL), often referred to as “good” and “bad” cholesterol.

The good carriers (HDL) bring cholesterol to the liver to be processed. **The higher the HDL level in your blood, the lower the risk of cardiovascular disease.**

Conversely, the bad carriers (LDL) make it easier for cholesterol to deposit on artery walls, which leads to atherosclerosis (hardening of the arteries). **The more LDL in the blood, the greater the risk of cardiovascular disease.**

Generally, the desirable total cholesterol level is under 5.2 mmol/L. A total cholesterol level that's lower or higher than 5.2 mmol/L may be considered a problem depending on the ratio of good (HDL) and bad (LDL) cholesterol as well as other risk factors.

There is also another type of fat found in the blood called **triglycerides**. They also increase the risk of cardiovascular disease, especially when other cholesterol levels are abnormal.



Eat healthy



COUNT ON “WINNING CHOICES”

Every day, ideally at every meal, choose “winning choices” among the four food groups. These foods are full of vitamins, minerals, antioxidants, fibres, and should be the basis of your diet.

Make more
space for them
in your diet



Vegetables and Fruit

EAT 7 TO 10 SERVINGS A DAY

Eat at least one dark green vegetable (broccoli, spinach, green peas, etc.) and one orange vegetable (carrots, sweet potatoes, etc.) every day.

Enjoy main courses filled with vegetables such as stew, couscous, stir-fry and salad.

You haven't had your fruit at mealtime? Have it for a snack!

Grain Products

OPT FOR WHOLE GRAIN PRODUCTS

Discover the many breads made from whole wheat and other grains, as well as whole wheat pasta, brown rice, oat or bran breakfast cereals, etc. They are an excellent source of energy and naturally low in fat.

Favour
lighter choices



Milk and Alternatives

CHOOSE MILK AND YOGURT THAT CONTAIN 2% OR LESS M.F.

You can also try calcium enriched soy beverages.

Cheese at 20% M.F. or less is a better option. Even at 20%, go easy if you have a cholesterol or a weight problem.

Meat and Alternatives

VARY YOUR SOURCES OF PROTEIN

Choose leaner cuts of red meat and poultry without the skin; go for small servings.

Have fish at least 2 or 3 times a week; don't shy away from fatty fish that contain good omega-3 fat.

Legumes... and even tofu can be interesting alternatives to meat.

As for unsaturated fat ...

Certain high-fat foods should be part of our diet because they are a source of unsaturated fat (mono and poly-unsaturated, including omega-3) whose health benefits have been widely recognized.

CHOOSE AMONG:

- vegetable oil such as canola, olive, nut, soybean, etc.
- non-hydrogenated soft margarine
- nuts and seeds
(don't forget ground flaxseed)
- avocado

CAUTION!

These foods are high in fat and also in calories. They should be eaten regularly but **in moderation**, especially if you have to watch your weight.

Eating healthy is a must to reduce the risk of cardiovascular disease, manage your weight, the levels of cholesterol and triglycerides in your blood, your blood pressure and diabetes. And don't forget that "eating well" is a source of energy and ... pleasure!

MANAGE THE FAT, THE SUGAR AND THE SALT

MANAGE DOES NOT MEAN ELIMINATE.

It's all in the frequency and the quantity!

↓ FAT

Reducing high-fat foods, especially those rich in **trans fat, saturated fat and cholesterol**, is highly recommended to individuals with elevated levels of cholesterol or triglycerides and to diabetics. Managing your fat intake also helps in controlling your weight.

Processed foods containing partially hydrogenated oil or shortening are major sources of trans fat. They are also rich in fat and calories:

- high-fat baked goods (pastries, croissants, pies, cakes, cookies, muffins)
- fried foods (French fries, chips, doughnuts, breaded foods)
- hydrogenated margarine, shortening

Fatty dairy products and meat are the major sources of saturated fat and cholesterol:

- butter, cream and ice cream
- high-fat cheese
- high-fat meat
- high-fat deli meats, sausages, bacon and pâté

If your blood cholesterol level is elevated, it is best you watch out for certain foods that are high sources of cholesterol:

- avoid fancy meats (liver, kidney, brain, sweetbread)
- stick to 2 to 3 egg yolks a week
- eat shrimp sparingly

↓ SUGAR

Reducing sweets is strongly recommended to diabetics and to individuals whose blood triglyceride level is elevated.

- sugar
- honey
- brown sugar
- syrup
- jams

These foods can add a sweet note to nutritious foods such as plain yogurt, milk-based or fruit desserts or home-made muffins but, it is best to limit the quantity.

- soft drinks
- fruit flavoured beverages

These beverages only offer sugar and calories, without nutritional value.

- pastries
- cakes
- pies
- cookies
- chocolate
- frozen desserts

These foods are often very sweet and very rich in bad quality fat and calories.

↓ SALT

Reducing the intake of salt and salty foods is recommended to individuals with high blood pressure.

More than 75% of the salt we eat is hidden in processed foods:

- fast-food
- ready-to-eat meals
- deli meats
- canned soups
- commercial sauces and broth
- chips, pretzels and other salty snack foods

Replacing salt

Herbs, pepper, lemon, garlic or ginger can add even more taste than salt.



LISTEN TO YOUR BODY!

Eating only when hungry and stopping when satisfied, are good ways to not overeat.

ONE OF 4 WINNING STRATEGIES



EAT HEALTHY

Questions of the

Alcohol

A moderate alcohol consumption can be beneficial to your heart. By moderate, we mean **no more than one or two drinks* per day**, for a maximum of 14 per week for men and 9 for women. If you don't drink or drink very little, don't start drinking to have a healthy heart. Individuals with high blood pressure or a high level of triglycerides in the blood should be more mindful of their alcohol consumption. Pregnant women, should avoid it altogether. As for those who are watching their weight, don't forget alcohol is an important source of calories.

* One drink =

- Regular beer (340 mL / 12 ounces)
- Regular wine (150 mL / 5 ounces)
- Spirits (45 mL / 1 1/2 ounces)

Stress


More and more, current research is showing a direct relationship between stress levels and cardiovascular disease. Stress can certainly have an impact on your heart's health when it leads you to eat poorly, smoke more often and neglect physical activity.

It's logical to think that depression, anxiety and the lack of support can also have a negative impact on your heart. These factors affect your ability to follow a treatment, or to acquire and maintain a healthy lifestyle. Without seeing them as the only factors responsible for a heart attack or stroke, these psychological factors deserve to be taken into consideration in the prevention and treatment of cardiovascular disease.

Women's heart

Many women don't consider cardiovascular disease as an important threat to their health. Yet cardiovascular disease is an "equal opportunity" disease, and **the main cause of death among both women and men**. In fact, cardiovascular disease is responsible for about 31% of deaths among men and 33% among women.

It is true that before menopause, women benefit from better natural protection than men. The risk increases gradually after menopause, however, because of reduced estrogen production. Generally, women are affected by cardiovascular disease 10 to 15 years later than men, the critical point being after age 65.



Medication, if necessary

For everyone, the vital first step in the battle against cardiovascular disease is to acquire and maintain healthy lifestyle habits. In some cases, however, it won't be enough to control an abnormal cholesterol level, elevated blood pressure or diabetes. Under these circumstances, **your physician will decide with you whether medication is an option in your case.**

As these conditions generally don't produce any symptoms or make people feel ill, 50% of patients stop taking their medication after a year and put their health – even their life – in danger. We now have very effective medication with far fewer side effects than in the past. If you experience side effects, your physician can usually adjust the dosage or suggest alternatives. It is important to **follow the prescription** and not change it without first talking with your physician or pharmacist.

WHAT ABOUT NATURAL PRODUCTS?

Just because something is natural, doesn't mean that it's not dangerous. Some products can cause side effects, interact with other medication or must be avoided in the case of certain illnesses. Unlike medication, these products are not subject to rigorous testing prior to being marketed. Before buying or taking natural

Weight chart

- 1 Find your height (in centimetres or feet and inches) in the blue column.
- 2 On the same line, find your current weight zone (in kilograms or pounds).

This weight chart is based on BMI* and is designed for adult men and women aged 20 to 65. It does not apply to children, adolescents or pregnant women. If you are very muscular or are over 65, your weight can slightly exceed the suggested limit without necessarily increasing your risk level.

* BMI = body mass index (kg/m²).

HEIGHT		WEIGHT							
cm	(ft.in.)	ZONE 1 Underweight BMI: under 18.5		ZONE 2 Normal weight BMI: 18.5 to 24		ZONE 3 Overweight BMI: 25 to 29		ZONE 4 Obesity BMI: 30 and over	
		under						over	
		kg	(lb)	kg	(lb)	kg	(lb)	kg	(lb)
145	4'9"	39	86	39-53	86-117	53-63	117-139	63	139
148	4'10"	41	89	41-55	89-121	55-66	121-146	66	146
150	4'11"	42	92	42-56	92-123	56-68	123-150	68	150
152	5'0"	43	94	43-58	94-128	58-69	128-152	69	152
155	5'1"	44	98	44-60	98-132	60-72	132-159	72	159
158	5'2"	46	102	46-62	102-137	62-75	137-165	75	165
160	5'3"	47	104	47-64	104-141	64-77	141-170	77	170
162	5'4"	49	107	49-66	107-146	66-79	146-174	79	174
165	5'5"	50	111	50-68	111-150	68-82	150-181	82	181
168	5'6"	52	115	52-71	115-156	71-85	156-187	85	187
170	5'7"	53	118	53-72	118-159	72-87	159-192	87	192
172	5'8"	55	120	55-74	120-163	74-89	163-196	89	196
175	5'9"	57	125	57-77	125-170	77-92	170-203	92	203
177	5'10"	58	128	58-78	128-172	78-94	172-207	94	207
180	5'11"	60	132	60-81	132-179	81-97	179-214	97	214
183	6'0"	62	136	62-84	136-185	84-100	185-221	100	221
185	6'1"	63	139	63-86	139-190	86-103	190-227	103	227
188	6'2"	65	144	65-88	144-194	88-106	194-234	106	234
190	6'3"	67	147	67-90	147-198	90-108	198-238	108	238
193	6'4"	69	152	69-93	152-205	93-112	205-247	112	247

Source: Health Canada. Canadian Guidelines for Body Weight Classification in Adults, 2003.

The **warning signs** of cardiovascular **disease**

To be able to act quickly when a heart attack or stroke occurs, you must know how to recognize the warning signs. Of course, you don't have to rush to the hospital at the slightest symptom, but if you have

one or more of the signs described below, go to the hospital emergency **QUICKLY**. Being admitted within a maximum of two hours following the appearance of symptoms could save your life.

ANGINA

This is a **partial blockage** of a heart artery that prevents blood from nourishing the part of the heart usually served by this artery. This can cause the following warning signs:

- different types of chest pain
- with or without pain in the arm, neck or jaw
- lasts a few seconds to a few minutes
- occurs most often with effort or strong emotions
- generally relieved within 5 or 10 minutes by rest or nitro-glycerine taken under the tongue or by spray.

WOMEN

For many women (and also certain men), angina can happen while at rest. Some may feel more pain in the neck or lower jaw, or they may feel out of breath, nauseated or tired instead of or in addition to having chest pain.

HEART ATTACK

This is a **total blockage** of a heart artery that completely prevents blood from nourishing the part of the heart usually served by this artery. This can cause the following warning signs:

- severe chest pain, taking different forms: a feeling of tightening, pressure, heaviness or burning, etc.
- may be accompanied by the following symptoms:
 - pain that spreads to shoulders, arms, neck, jaw or back
 - dizziness, paleness, difficulty breathing, sweating, nausea, vomiting or anxiety
- usually lasts more than 10 minutes
- not relieved by rest or by 2 or 3 doses of nitro-glycerine
- can sometimes occur silently, with none of the above symptoms.

WOMEN

For many women (and also for certain men), this classic list of symptoms is replaced by more vague symptoms:

- pain in the neck, jaw or arms
- heaviness in the shoulder blades, back or the pit of the stomach
- feeling out of breath, tired, sweating, nausea, vomiting.

An estimated 50% of heart attacks in women cause no severe pain.

STROKE

This is a blockage of an artery in the neck or head that prevents blood from nourishing the part of the brain usually served by this artery. This can cause the following warning signs:

- sudden weakness or loss of sensation of the mouth, arm or leg, on only one side of the body
- sudden reduction or loss of vision
- difficulty or inability to speak
- dizziness or loss of balance.

Note that these symptoms can sometimes be more vague, particularly among the elderly. Watch for a sudden change in behaviour, like unusual slowness, indifference or an inability to dress or feed oneself.

YOU COULD SAVE A LIFE

Rapid intervention to assist a victim of a heart attack – within the first few minutes – can save a life. To learn how to do cardiopulmonary resuscitation (CPR), contact the Heart and Stroke Foundation toll free at: 1 888 473-4636.

Certain symptoms listed in these descriptions may not be linked to a cardiovascular problem. When in doubt, it is better to let the physician be the judge rather than to diagnose oneself.

The original version of this document was developed by Doctors Louis Gagnon and Gilles Pineau and by dietitians Sylvie Desroches and Danièle Prévost of the ACTI-MENU Health Program as well as Raymonde Pineau, M.D., Direction de santé publique de la Capitale-Nationale and Élise Latour, Dietitian, Montreal Heart Institute, in collaboration with: Louise Labrie, Health promotion specialist and Isabelle Samson, M.D., Direction de santé publique de Montréal; Serge Goulet, M.D. and Huguette Bélanger, M.D., Direction de santé publique de la Montérégie; Roxane Néron, M.D., Direction de santé publique des Laurentides; François Croteau, M.D., Santa Cabrini Hospital; Brigitte Lachance, Ministère de la Santé et des Services sociaux du Québec, with the support of a scientific committee composed of: Renaldo Battista, Professor, Faculty of Medicine, University of Montreal; Monique D. Gélinas, Ph.D., Nutrition, University of Montreal; Jean Davignon, Internist, Clinical Research Institute of Montreal; Mireille Dubost, Dietitian, University of Montreal; Gaston Godin, Ph.D., Behavioural Sciences, Laval University; Martin Juneau, Cardiologist, Montreal Heart Institute; Réal Lacombe, M.D., Direction de santé publique de l'Abitibi-Témiscamingue; and André Lacroix, Endocrinologist, Centre hospitalier de l'Université de Montréal.

We would also like to thank the following experts who were consulted: Beth Abramson, Internist and Cardiologist, St Michael's Hospital, Toronto; Philip Barter, Cardiologist, Heart Research Institute, Sydney, Australia; Greg Bondy and Jiri Frolich, Medical Biochemists, St Paul's Hospital, Vancouver; W. Virgil Brown, Cardiologist, VAMC, Georgia, United-States; Jean-Pierre Després, Ph.D., Exercise Physiology, Institut de cardiologie de Québec; Robert Dufour, M.D., Clinical Research Institute of Montreal; George Fodor and Ruth McPherson, Internists, University of Ottawa Heart Institute; Jacques Genest and Allan Sniderman, Cardiologists, Royal Victoria Hospital, Montréal; Rafik Habib, M.D., Centre cardio-vasculaire de Laval; Pavel Hamet, Endocrinologist, Centre hospitalier de l'Université de Montréal; Marc-André Lavoie, Cardiologist, Montreal Heart Institute; Gilles Paradis, Epidemiologist, Direction de santé publique de Montréal; Barry Rose, M.D., F.R.C.P(C), F.A.C.C., Health Sciences Complex of St. John's Newfoundland; George Steiner, Endocrinologist, Toronto General Hospital; James Stone, Cardiologist, University of Calgary; Guy Tremblay, Cardiologist, Hôpital du Saint-Sacrement, Québec. Graphic Art: M. Edgar Inc.

Professionals and companies may order this brochure by calling (514) 326-2466 or 1-866-356-2466.

© ACTI-MENU - 1993 for the first publication

© ACTI-MENU - 2007 for this publication. Commercial use is forbidden. Any reproduction is prohibited without written authorization.

Visit our website at www.actimenu.ca

Legal Deposit - Bibliothèque et Archives nationales du Québec, 2007.